

National
Library
of MedicineMy NC
[Sign In] [Regis

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search PubMed for Display Abstract 20 All: 1 Review: 0 [About Entrez](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)[E-Utilities](#)[PubMed Services](#)[Journals Database](#)[MeSH Database](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[LinkOut](#)[My NCBI \(Cubby\)](#)[Related Resources](#)[Order Documents](#)[NLM Catalog](#)[NLM Gateway](#)[TOXNET](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[PubMed Central](#)☐ 1: Mol Endocrinol. 1988 Mar;2(3):277-83.[Related Articles, Links](#)

A Moloney MLV-rat somatotropin fusion gene produces biologically active somatotropin in a transgenic pig.

Ebert KM, Low MJ, Overstrom EW, Buonomo FC, Baile CA, Roberts TM, Lee A, Mandel G, Goodman RH.

Department of Anatomy and Cellular Biology, Tufts University School of Veterinary Medicine, Grafton, Massachusetts 01536.

Expression of a Moloney murine leukemia virus (MLV) rat somatotropin fusion gene was examined in a transgenic pig. The fusion gene was integrated in a single site within the genome in a tandem array with approximately eight copies per cell. The integrated in a single site within the genome in a tandem array with approximately eight copies per cell. The integrated MLV-rat somatotropin fusion gene produced high levels of circulating rat somatotropin and resulted in an elevation in the circulating levels of insulin-like growth factor I. Although there was no increase in the rate of growth of the transgenic animal during the rapid growth phase, several phenotypic changes were evident. Skeletal growth was markedly increased and fat deposition was reduced throughout the animal. Blood glucose levels were elevated without ketosis. Northern blot analyses of rat somatotropin RNA revealed that expression of the fusion gene was highest in the spleen, lung, intestine, lymph nodes, and bone marrow. These results show that the MLV promoter can be used to express high levels of biologically active rat somatotropin in transgenic swine.

PMID: 3398854 [PubMed - indexed for MEDLINE]

Display Abstract 20 [Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)